

**Screening, Brief
Intervention and Referral to Treatment (SBIRT)
in Behavioral Healthcare**

I. INTRODUCTION

This report discusses the evidence supporting the effectiveness of screening, brief intervention, and referral to treatment (SBIRT) as a comprehensive approach, as well as for the implementation and effectiveness of the individual components of SBIRT for different behavioral health conditions.¹ The report describes briefly the underlying research that has been conducted in the prevention and early intervention of risky alcohol, substance abuse and tobacco consumption, as well as commonly reported mental health problems, and describes existing studies/models for specific populations and settings. Further, the report addresses the question of what a model SBIRT program is, compared to programs which include or adapt components of the comprehensive SBIRT approach. Literature reviews are included in Attachment I. This paper is intended for use by policy makers, research organizations and governmental agencies seeking to understand the complexities of the SBIRT model and/or considering the adoption and implementation of SBIRT systems change or behavioral health integration within primary care settings.

Screening, brief intervention, and referral to treatment (SBIRT) was originally developed as a public health model designed to provide universal screening, secondary prevention² (detecting risky or hazardous substance use before the onset of abuse or dependence), early intervention, and treatment for people who have problematic or hazardous alcohol problems within primary care and other health care settings (Babor et al., 2007; Babor & Higgins-Biddle, 2001). Based on the SAMHSA model, SBIRT is unique in its universal screening of all patients regardless of an identified disorder, allowing health care professionals to address the spectrum of such behavioral health problems even when the patient is not actively seeking an intervention or treatment for his or her problems.

Following are the key points of this paper:

- SBIRT has been defined by SAMHSA as a comprehensive, integrated, public health approach to the delivery of early intervention for individuals with risky alcohol and drug use, and the timely referral to more intensive substance abuse treatment for those who have substance abuse disorders. There is consensus that a comprehensive SBIRT model includes screening, brief intervention/brief treatment and referral to treatment. In addition to these

¹ Excludes medical conditions.

² There is some discussion about whether SBIRT is selective prevention (Kumpfer & Baxley, 1997) or early intervention given the overlap in SBIRT's approach and objectives.

integral components, SAMHSA defines a comprehensive SBIRT model to include the following characteristics:

- It is brief (e.g., typically about 5-10 minutes for brief interventions; about 5 to 12 sessions for brief treatments).
 - The screening is universal.
 - One or more specific behaviors related to risky alcohol and drug use are targeted.
 - The services occur in a public health non-substance abuse treatment setting.
 - It is comprehensive (comprised of screening, brief intervention/treatment, and referral to treatment).
 - Strong research or experiential evidence supports the model's effectiveness.
- No standard SBIRT definition has been articulated by the U.S. Preventive Services Task Force or other authoritative/coordinating bodies. The SAMHSA definition of SBIRT is based on methodology that was developed during the implementation of a comprehensive SBIRT grant program comprised of all the integral components, and supported by research by the National Institute on Drug Abuse and the National Institute on Alcohol Abuse and Alcoholism.
- There is substantial research on the effectiveness of SBIRT in reducing risky alcohol consumption. However, the evidence for the effectiveness of SBIRT in reducing risky drug use, although promising, is still accumulating. The results for the SAMHSA model of SBIRT for drug misuse are inconsistent depending on the characteristics of the provider, the specific setting, and the patient population that is targeted for SBIRT implementation. While there is robust evidence for screening and referral for depression in primary care, to date, little empirical evidence for the use of comprehensive SBIRT-like models for mental health problems commonly reported by health care patients. There is also no research that has demonstrated the implementation or effectiveness of SBIRT-like models in addressing trauma or anxiety disorders in clinical health settings.

II. THE SAMHSA SBIRT MODEL

SBIRT is a comprehensive, integrated, public health approach to the delivery of early intervention for individuals with risky alcohol and drug use, as well as the timely referral to more intensive substance abuse treatment for those who have substance use disorders. Primary care centers, hospital emergency rooms, trauma centers, and community health settings provide opportunities for early intervention with at-risk substance users before more severe consequences occur.

SAMHSA supports a research based comprehensive behavioral health SBIRT model which reflects the six following characteristics:

1. It is brief. The initial screening is accomplished quickly (modal time about 5-10 minutes) and the intervention and treatment components indicated by the screening results are completed in significantly less time than traditional substance abuse specialty care.

2. The screening is universal. The patients, clients, students, or other target populations are all screened as part of the standard intake process.
3. One or more specific behaviors are targeted. The screening tool addresses a specific behavioral characteristic deemed to be problematic, or pre-conditional to substance dependence or other diagnoses.
4. The services occur in a public health, or other non-substance abuse treatment setting. This may be an emergency department, primary care physician's office, school, etc.
5. It is comprehensive. The program includes a seamless transition between brief universal screening, a brief intervention and/or brief treatment, and referral to specialty substance abuse care.
6. Strong research or substantial experiential evidence supports the model. At a minimum, programmatic outcomes demonstrate a successful approach.

As a comprehensive or model approach, SBIRT has only been demonstrated to be effective for risky alcohol use. There is substantial evidence for the effectiveness of brief interventions for harmful drinking when delivered by a physician or other qualified health professional (Bien et al, 1993; Kahan et al, 1995; Wilk et al, 1993). There is a growing body of literature showing the effectiveness of SBIRT for risky drug use (Madras et al, 2008; Saitz et al, 2010; Bernstein et al., 2005) but the results vary by the characteristics of the provider, the specific setting, and the patient population that is targeted for SBIRT implementation.

To determine the effectiveness of SBIRT beyond alcohol, a comprehensive literature review was conducted. SBIRT-like models including not only a simple screening tool, but also an appropriate and brief intervention that addressed the level of problem indicated by the screening results. Table 1 (p. 4) identifies the substance abuse and mental health conditions where SBIRT or components of SBIRT have been employed. The literature review did not include studies that employed SBIRT or approaches that are similar to SBIRT for general medical conditions such as blood pressure, HIV/AIDS, or other behavioral issues such as domestic violence.

As shown in Table 1, the comprehensive SBIRT model has not been consistently demonstrated as effective in addressing harmful or risky drug misuse, depression, trauma, or anxiety problems. Findings showing the effectiveness of SBIRT for drug misuse are accumulating, and there is some programmatic data from the SAMHSA State SBIRT programs showing promising findings for depression among primary care patients. Public health approaches that are consistent with the SBIRT model have also been demonstrated for tobacco use. They are described in the latter sections of this paper. Table 1 presents a brief analysis of the evidence for the effectiveness of SBIRT for various behavioral health conditions.

Table 1. EFFECTIVENESS OF SBIRT AND ITS COMPONENTS FOR BEHAVIORAL HEALTH CONDITIONS

	Screening	Brief Intervention ¹	Brief Treatment ²	Referral to Treatment	Evidence for Effectiveness of SBIRT
Alcohol Misuse/Abuse	✓	✓	✓	✓	Comprehensive SBIRT effective (Category B classification, USPSTF)
Illicit Drug Misuse/Abuse	✓	*	*	✓	Growing but inconsistent evidence
Tobacco Use	✓	✓	✓	✓	Effective brief approach consistent with SBIRT (USPSTF; 2008 U.S. Public Health Service (PHS) Clinical Practice Guideline)
Depression	✓	—	✓	✓	No evidence to date for depression
Trauma/Anxiety Disorders	✓	*	—	✓	No evidence to date for trauma/anxiety disorders


Key: ✓ Evidence for effectiveness/utility of component

* Component Demonstrated to show Promising Results

— Not Demonstrated and/or Not Utilized

¹Brief intervention as defined by the SAMHSA SBIRT program involves 1-5 sessions lasting 5 minutes to an hour. Among SBIRT grantees funded by SAMHSA, about 15% of patients receive scores that indicate a brief intervention.

²Brief treatment as part of SBIRT involves 5-12 sessions, lasting up to an hour. Among State SBIRT grantees funded by SAMHSA, about 3% of patients receive a score that dictates a brief treatment.

Chart 1. FLOW CHART FOR SBIRT PROCESS

Screening

Universal screening helps identify the appropriate level of services needed based on the patient's risk level. Patients who indicate little or no risky behavior and have a low screening score may not need an intervention. Those who have moderate risky behaviors and/or reach a moderate threshold on the screening instrument may be referred to brief intervention. Patients who score high may need either a brief treatment or further diagnostic assessment and more intensive, long term specialty treatment. Screening typically takes 5-10 minutes and can be repeated at various intervals as needed to determine changes in patients' progress over time. Some commonly used screens for the implementation of SBIRT for alcohol and drug use are the Alcohol Use Disorders Identification Test (AUDIT), Drug Abuse Screening Test (DAST), Alcohol, Smoking, Substance Involvement, Screening Test (ASSIST), and the Cut Down, Annoyed, Guilty, Eye-Opener (CAGE). In addition, a recent study found a single question related to drug use to be effective in detecting drug use among primary care patients (Smith et al., 2010).

Prescreening, which is not a core component of SBIRT but is frequently used, reduces the time needed by busy clinic staff to identify patients with risky behavior. Examples of validated pre-screens are the Alcohol Use Disorders Identification Test-Consumption (AUDIT-C), which consists of the first three alcohol consumption questions from the full 10-item AUDIT questionnaire, and the NIAAA prescreening question ("On any single occasion during the past 3 months, have you had more than 5 drinks containing alcohol?", Taj et al., 1998). If a patient scores high on any domain in the pre-screen, a full screen is conducted.

Brief Intervention (BI) and/or Brief Treatment

Patients are provided with BI, brief treatment, or referral to intensive specialty treatment depending on their level of risk using a validated pre-screen and/or screening tool (Babor & Higgins-Biddle, 2001). With respect to substance abuse, in general only a small proportion of patients in primary care settings screened positive for some level of substance misuse, abuse or dependency. This is usually 5%-20%, but may be as high as 40% in some clinical settings. The majority of patients report minimal or no problems with alcohol or drugs and as such may be an ideal group for primary or universal prevention activities for maintenance of non-risky use or abstinence. The goal of a BI (which usually involves 1-5 sessions lasting about 5 minutes to one hour) is to educate patients and increase their motivation to reduce risky behavior.

The goal of brief treatment (which usually involves 5-12 sessions) is to change not only the immediate behavior or thoughts about a risky behavior but also to address long-standing problems with harmful drinking and drug misuse and help patients with higher levels of disorder obtain more long term care. Based on performance data from state SBIRT grantees funded by SAMHSA, only about 3% receive a score that indicates a brief treatment. Patients referred to a brief treatment often have higher risk factors than those referred to a BI. Brief treatment may also require a manualized course of (advanced) motivational enhancement and cognitive behavioral approaches to help patients address unhealthy cognitions and behaviors associated with current use patterns and adopt change strategies. If patients report greater risk factors than what brief treatment can address, they are referred to specialty substance abuse care. In some cases, a patient may receive a BI first and then move on to a brief treatment or longer term care. Although the time required to execute BI/BT is generally considered brief, it is far too lengthy for physicians to do. Also, physicians cite concerns about angering or insulting patients by bringing up sensitive issues such as alcohol and/or drug use. While these concerns are understandable, when SBIRT is implemented properly, the time commitment is reasonable and acceptably low given the demonstrated success in identifying persons requiring referral to treatment (RT). Similarly, concerns about patient reactions can be neutralized by proper training for the providers and ensuring that access to referral services is available. In addition, SBIRT is frequently implemented by allied health professionals such as nurses, social workers, or health educators, with results and actions noted in the patient chart for physician notification and oversight.

Referral to Treatment (RT)

Referral to treatment can be a complex process involving coordination across different types of services. As such, the absence of linkages to treatment referrals can be a significant barrier to the adoption of SBIRT. Referral is recommended when patients meet the diagnostic criteria for substance dependence or other mental illnesses as defined by the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV).³ In these cases, a referral to a specialized treatment provider is often made. Referral requires the primary care system to establish new and complex linkages with the traditional specialty care system to connect clients who score in the problematic range to recognized, evidence based treatment in a timely manner. Although only 3% to 4% of screened patients in primary care settings typically need to be referred, the absence of a proper treatment referral will prevent the patient from accessing appropriate and timely care that can impact other psychosocial and medical issues. Research findings suggest that motivational-based BIs can increase patient participation and retention in substance abuse treatment (Hillman et al., 2001; Dunn and Ries, 1997). Strong referral linkages are critical, as well as tracking patient referrals. SAMHSA requires SBIRT grantees to have a comprehensive referral to treatment and follow-up system in place for the duration of the program. In the case where RT is incorporated into an integrated care model, this may require shifts in provider allocation and hiring.

³ The diagnostic criteria are likely to change when DSM V is released in 2012 or 2013.

The following characteristics of SBIRT identified in the research literature (see Reference section) have been shown to be important in effectively addressing behavioral health problems. They have therefore formed the foundation for the SAMHSA SBIRT programs.

- 1) Use of brief, validated, universal pre-screening/screening tools. These tools allow health care professionals to address the problem behavior even when the patient is not actively seeking treatment for his or her problem. Prescreening/screening tools accurately and quickly identify individuals with problematic conditions in as little time as 2-4 minutes. Because of its brevity and its universal application (that is, can be used with all patients), SBIRT may be more generally accepted by health care professionals working in busy practices.
- 2) Relatively easy to learn by diverse providers. The SBIRT approach is easy to learn relative to other behavioral treatment techniques that may require lengthy specialized training. As such, it can be implemented by diverse health professionals who work in busy medical settings such as physicians, nurses, social workers, health educators and paraprofessionals.
- 3) Incorporation of strong referral linkages to specialty treatment. Approaches that are effective integrate comprehensive strategies that include referral to specialty treatments (Gentilelo, Donovan, Dunn & Rivara, 1999). While RT may be difficult in underserved areas, this should not deter programs from engaging in developing SBI activities as they have beneficial effects separate from the referral. However, the goal is to provide a quick handoff for dependent patients to specialty treatment if the primary care site cannot provide more intensive services for substance abuse. Establishing linkages with specialty care through identification of local treatment service contracts, an MOU agreement between sites, or dedicated central referral services has been a major barrier for many providers in their decision to adopt SBIRT. The availability of well established referral linkages to specialty care is essential to the uptake and maintenance of SBIRT, and closely tracking to confirm patient compliance with treatment is critical to good health care provision. Primary care locations engaged in referral to specialty care make efforts to determine the patient's engagement and participation in treatment, as this may also affect the course of treatment in the general medical practice.

III. ALCOHOL MISUSE, ABUSE, AND PREVENTION

There is substantial evidence from review studies (Babor, 2007; Bein et al, 1993; Kaner, et al., 2009) and meta-analyses of randomized clinical trials (Beich et al., 2003; Bertholet et al, 2005) that show the effectiveness of SBIRT in reducing hazardous drinking in patients presenting in primary care and other health care settings. The U.S. Preventative Services Task Force (USPSTF) has recommended that “behavioral counseling interventions for risky/harmful alcohol use among adult primary care patients can provide an effective public health approach to reducing problematic drinking” (USPSTF, 2004). The USPSTF also concluded that counseling for risky drinkers should include advice to reduce current drinking; feedback about current drinking patterns; and explicit goal-setting, usually for moderation and assistance in achieving the goals.

Research also indicates that despite the robustness of the evidence for SBIRT's effectiveness for unhealthy alcohol drinking, other factors can impact its effects. For example, studies have shown that multiple contacts or sessions (in contrast to a single contact) with a provider can increase the impact of SBIRT in reducing risky alcohol consumption (Brown et al., 2007; Longabaugh et al., 2001). Moreover, demographic factors and psychosocial conditions also have been shown to influence SBIRT's effects on alcohol misuse (Saitz et al., 2006). For example, homelessness makes SBIRT less effective due to the challenges involved in working with this population, and brief interventions have improved linkages with those who can provide assistance to younger men and hospitalized women.

The conduct of universal screening, brief intervention and treatment, and referral to treatment for alcohol disorders has been found to be effective in various healthcare settings for diverse patient populations including primary care (Babor et al., 2007), emergency departments (Gentilello et al., 1999), as well as schools and colleges (O'Brian et al., 2006). Data are currently being collected that suggest that SBIRT may also be effective in addressing alcohol problems in employee assistance programs (McPherson and Goplerud, 2008). Recent research also has demonstrated the efficacy of conducting screenings and BIs using innovative strategies such as the use of personalized feedback via the internet (Cunningham, 2010), as well as web-based outcomes monitoring to assist with treatment decisions and cognitive behavioral techniques (Roy-Bryne, 2010).

Also promising is the utilization of computerized interventions which has been shown to be effective in augmenting and complementing the gains made through the initial face to face brief interventions. The Veterans Administration, for example, examined the use of electronic clinical reminders with patients following screening with the AUDIT-C and showed such approaches reinforced moderate drinking reductions at follow up (Williams, 2010). Other research reviews indicate that electronic methods can enhance brief interventions with substance users by offering assessment and feedback in brief motivational interviewing; monitoring individual treatment patient's progress; tracking patients in aftercare; and providing educational opportunities for clinicians (Cucciare, 2009). Electronic intervention can also help bridge the treatment capacity gap by providing another source of assistance for women who do not complete traditional substance abuse treatment (Van DeMark, et al., 2010). In addition, the cost savings offered by the implementation of SBIRT in primary care are significant. One study (Gentilello, 2005) showed that for every one dollar spent on providing SBIRT approximately \$3.81 is saved. The Washington State SBIRT program cost study also reflects similar savings.

The concept of SBIRT can be applied across the continuum of care for alcohol problems. Based on the severity of the problem indicated by the screening results, interventions ranging from universal prevention to brief interventions to traditional specialty treatment can be provided to health care patients. For individuals who are abstinent, universal prevention practices can be implemented to sustain alcohol abstinence. For moderate risky drinking, the first two components of SBIRT – screening and brief interventions (SBI)– may be implemented which can address inappropriate expectancies (beliefs about substance use effects and social norms of acceptable behavior) and lack of motivation to change risk factors that contribute to substance abuse (Dimeff et al., 1999).

Extensive research supports screening and brief intervention as effective universal and selective prevention strategies for alcohol problems. Universal screening with educational content has measurable prevention effects with accompanying feedback (Kunz et al., 2004). The prevention approach may also be successful for abstainers and non-risky drinkers by providing behavioral support and normative information to maintain healthy behaviors. For at-risk individuals, early identification and brief intervention around false expectancies, normative use misperceptions and skills acquisition can prevent progression to severe drinking problems. For example, the BASICS program, which is consistent with the SBIRT approach, has been shown to be effective in addressing problematic or risky drinking in college age groups (Dimeff et al., 1999). SBIs also incorporate motivational interviewing components (Miller and Rollnick, 2002) that are also integrated in brief treatment for higher risk patients. SBIs have proven effective in decreasing overall consumption and binge drinking (Casset et al., 2008; Hanewinkel & Wiborg 2005; Kunz Jr. et al., 2004; Martens et al., 2007; Heather et al., 2004; Toumbourou et al. 2007; Murphy et al., 2001), as well as increasing productivity (Osilla et al., 2010). Evidence further demonstrates that strengthening resiliency, competencies, and social connectedness supports recovery for those individuals who show early symptoms of alcohol misuse.

Extensive reviews of the effectiveness of SBI (Babor et al., 2007, 2008) have found that there are “irrefutable” improvements in short-term health benefits as well as indications of “substantial” long-term benefits. Follow up at three, six or nine month intervals can help document the effectiveness of SBI and reinforces normative ideation and skills enhancement for individuals with minimal risk behaviors. To achieve long term effects, SBI must be implemented with fidelity through targeted training for providers (Cameron et al., 2010; Seale et al., 2005; Christensen et al., 2004; Bray et al., 2009; Ronzani et al., 2008; Furtado et al., 2008; Heather et al., 2004; Tollison et al., 2008; Babor et al., 2004; Brown & Fleming, 1998). In many instances providers implementing SBI may not necessarily be physicians but allied health professionals such as nurses, counselors, health educators, and peers (Mastroleo, 2009; Blume & Marlatt, 2004), who may experience fewer barriers in service provision than physicians (Babor et al., 2004). Also, SBI can be conducted individually or with groups (Shellenberger et al., 2009; Henslee, 2009), with web-based instruments (i.e. college oriented E-Chug and E-Toke or Alcohol Skills Training Programs), or online feedback (Blume & Marlatt, 2004), and applied through strategic planning by communities or providers.

IV. DRUG MISUSE, ABUSE, AND PREVENTION

In 1995, based on the scant availability of published research on SBIRT for drugs, the USPSTF (1995) determined that there was “insufficient evidence to recommend for or against” the effectiveness of using an SBIRT approach for drugs. Some researchers have cited the relative scarcity of validated brief drug screening tools (Smith PC, et. al., 2010) and the low prevalence rates of drug use (Saitz, 2010) in primary care settings, as two reasons for the comparatively small number of studies showing SBIRT’s effects with drugs (De Micheli D, et. al., 2004). Nevertheless, since 1995, there has been a growing body of investigator-initiated research as well as findings from SAMHSA-funded SBIRT projects that have shown promising results for the use of the comprehensive SBIRT approach, as well as selected use of individual components, in reducing risky drug use (Copeland et al., 2001). For instance, a randomized controlled trial indicated that BIs can reduce cocaine and heroin use (Bernstein et al., 2005). Motivational

interviewing coupled with a self-help booklet given to regular amphetamine users also resulted in reduced levels of drug use (Baker, Lee, Claire, Lewin, Grant, & Pohlman, 2005). BIs for patients screening positive for cocaine, heroin, and amphetamine are also showing promising results in various settings beyond emergency departments (Cunningham et al., 2009). In small sample sizes, screening and BIs have been linked with reductions in the use of marijuana, amphetamine-type stimulants, cocaine, and heroin (Madras et al., 2008). The World Health Organization (2008) sponsored a multi-national study demonstrating that screening and brief interventions resulted in short-term reductions of a wide variety of illicit drugs, including marijuana, cocaine, amphetamine-type stimulants, and opioids.

As with alcohol consumption, universal and selective prevention efforts may also be targeted to those with minimal or mild drug misuse. Like with alcohol, identified abstainers can benefit from supportive and normative information to maintain healthy lifestyles. For individuals at risk for drug problems, early identification and brief intervention around false expectancies and skill acquisition can prevent progression to more severe drug problems. In addition, tools that can be used for universal screening of drug use in health settings such as the DAST and the ASSIST as well as on-line tools such as E-TOKE (Electronic – THC Online Knowledge Experience) are prevention-ready applications designed to detect the presence of drug use.

V. SBIRT AND TOBACCO USE

The utility of SBIRT approaches for all forms of tobacco use, especially smoking, has been endorsed by the USPSTF and has elicited interest in primary care and hospital personnel. Cigarette smoking continues to be the leading cause of preventable disease and death in the United States (USDHHS, 2004) and is attributed to approximately 443,000 deaths per year (CDCP, 2010) from lung cancer: ischemic heart disease, chronic obstructive pulmonary disease, strokes, and other diagnoses. Smoking also affects health outcomes of people other than the smokers, with smoking during pregnancy resulting in premature births, spontaneous abortions, stillbirths, and intrauterine growth retardation. In addition, research has shown that psychiatric disorders and cigarette smoking are frequently co-morbid conditions (Dome et al, 2010; Brown et al, 2008; Brown et al, 2002; Degenhardt & Hall, 2001; Grant et al, 2004). A recent study using data from the 2005-2006 National Survey on Drug Use and Health reported that adults with lifetime depression, anxiety, anxiety with depression, or major depressive episodes were more likely to be “current smokers, smoke with higher intensity and frequency, have more dependence, and have lower success at quitting” when compared to individuals without these psychiatric conditions (Troscclair & Dube, 2010).

However, despite smoking's established risks and the health benefits of quitting, 23 percent of adults in the United States continue to smoke and more than 2,000 adolescents become regular tobacco users daily (NSDUH, SAMHSA). Nearly 90 percent of smokers start by age 18, and 25 percent of teen smokers remain addicted as adults. Because 70 percent of smokers see a physician each year (Fiore, Bailey, Cohen, et al., 2000) clinicians have a unique opportunity to intervene and implement tobacco SBIRT in primary care settings and emergency departments.

As such, the USPSTF strongly recommends that clinicians screen all adults for tobacco use and provide brief interventions, including screening, brief behavioral counseling (less than 3 minutes), and pharmacotherapy delivered in primary care settings. The USPSTF also strongly

recommends that clinicians screen all pregnant women for tobacco use and provide augmented pregnancy-tailored counseling to those who use tobacco products. These interventions have been shown to be effective in increasing the proportion of smokers who successfully quit smoking and remain abstinent after 1 year.

The USPSTF advises that the clinical interventions for tobacco cessation that are cited in the 2008 U.S. Public Health Service (PHS) Clinical Practice Guideline, *Treating Tobacco Use and Dependence* (Fiore et al, 2008), become integrated in standard clinical practice. The PHS Guideline also recommends that clinicians use the screening instrument known as the 5A's of tobacco use intervention, which provides a useful strategy for engaging all medical patients in smoking cessation discussions. The 5A's are consistent with the SBIRT approach and parallel the screening and brief intervention or counseling components of the SBIRT model.

1. *Ask* about tobacco use.
2. *Advise* to quit through clear personalized messages.
3. *Assess* willingness to quit.
4. *Assist* to quit.
5. *Arrange* follow-up and support.

The Guideline's behavioral treatments include counseling, social support, problem solving, and cessation skills training offered in face-to-face individual or group formats or via telephone quit lines. Medication assisted treatments for tobacco use/dependence have also been suggested and include seven FDA-approved, first-line medications (i.e., bupropion SR, nicotine gum, inhaler, lozenge, nasal spray, and patch), and two second-line medications (clonidine and nortriptyline).

The Agency for Healthcare Research and Quality (AHRQ) also reviewed tobacco guidelines developed in England in 2006 and supports recommendations for brief interventions for patients who use tobacco products, including: simple advice to stop, assessment of the patients' commitment to quit, an offer of pharmacological or behavioral support, and provision of self help materials or referral to supportive resources such as Quit lines.

VI. DEPRESSION

The USPSTF supports screening for adult depression where accurate diagnosis, effective treatment, and follow-up are available. The USPSTF also recommends screening adolescents (12-18 years of age) for major depressive disorder (MDD), again with accurate diagnosis, psychotherapy (cognitive behavioral or interpersonal), and follow-up. There are many commonly used screening tools for depressive symptoms, such as the Patient Health Questionnaire 2 (PHQ-2) (Kroenke, et al., 2003) and the Patient Health Questionnaire 9 (PHQ-9) (Kroenke, et al., 2001) which both have established validity and reliability.

Primary care physicians are the providers most likely to see patients when they first become depressed and are most capable of initiating and monitoring treatments with pharmacologic agents (McNaughton, 2009). Previous studies, however, have shown that at least half of patients with active depression seen by primary care physicians remain undiagnosed (Spitzer et al, 1994; Schulberg et al., 1988; Ormell et al, (1991). Depression is particularly prevalent among "high

utilizers” of medical care resources, of whom as many as 40% have been found to have a current depressive illness (Katon et.al., 1990). Due to time constraints and training issues, physicians in primary care are often unable to provide effective behavioral interventions and treatments for the patients with mental disorders (McNaughton, 2009).

Promising but preliminary data are available from SBIRT grantees funded by SAMHSA that indicate that the SBIRT approach may be adapted for depression treatment. For example, the State of Wisconsin incorporated depression screening into a Wisconsin Initiative to Promote Healthy Lifestyles (WIPHL) pilot program. Patients with mild or moderate depression were provided behavioral activation by health educators using specific protocols developed by the program.

Behavioral activation also offers promise as a strategy for brief intervention and there is some evidence that it would fit an SBIRT-like approach. Behavioral activation assists individuals to identify and engage in daily activities and situations they find positively reinforcing and consistent with their long-term goals (Dimidjain et al., 2006). Behavioral activation as a brief intervention has been demonstrated in three meta-analyses, one randomized control trial, and one follow-up study of a previous randomized control trial, to be an effective intervention for the treatment of depression (Sturmey, 2009).

VII. ANXIETY DISORDERS AND TRAUMA

Anxiety disorders are among the most common mental health problems seen in primary care settings and as many as one-third of primary care patients have been found to have significant anxiety symptoms (Fifer, 1994). Approximately 15% of primary care patients have a current anxiety disorder, and 24% have had a lifetime anxiety disorder, as assessed by diagnostic interviews (Nisenson et al., 1998). Primary care patients with anxiety disorders typically have considerable disability and impairment in functioning (Roy-Byrne et al., 1999; Sherbourne et al., 1996) and high utilization rates of general medical services which ultimately result in higher health care costs (Simon et al., 1995). Screening tools are also available for anxiety such as the Brief Symptom Checklist-18 (Derogatis, 2001) which provides a measure of both anxiety and depression. The My Mood Monitor (M-3) (Gaynes et al., 2010) screening is a valid and efficient one page tool for screening multiple common psychiatric illnesses in primary care and other settings. The M-3 can function both as a screen for specific anxiety and mood disorder diagnoses, as well as a general screen for the presence of any mood or anxiety disorder in addition to bipolar disorder and PTSD.

Interventions such as passive psychoeducation, including bibliotherapy, have been shown to reduce symptoms of anxiety, psychological distress, and depression (Donker et al., 2009). These approaches may be offered as a brief intervention to patients who screen positive for mild or moderate levels of anxiety. Passive psychoeducational interventions are cost-effective and can be easily put into practice by non medical professionals and may have a less-stigmatizing impact on consumers, especially when delivered through a Web site, e-mail or a brochure (Donker et al., 2009).

Evidence of emotional trauma is also common in primary care. Walker et al.(1993) report that rates as high as 37% for childhood sexual abuse and 29% for adult sexual assault are evident in primary care settings. Walker et al. found that 61% of women reported that they believed that it was appropriate for their primary care physician to ask about previous victimization, but only 4 percent had been actually asked. In the Adverse Childhood Experiences (ACE) Study (Dube et al., 2004), patients received an assessment using the Family Health History and Health Appraisal questionnaires as measures. The authors found the reliability statistics of the ACE study support the use of these questionnaires for retrospective reports of adverse childhood experiences such as childhood maltreatment, household dysfunction, and other socio-behavioral factors. Other tools for screening trauma and anxiety include: the Trauma Symptom Inventory (Briere, 1995), the PTSD-8 (Hansen, et al., 2010), and the Primary Care PTSD Screen (PC-PTSD) (Prins, et al., 2003).

The National Child Traumatic Stress Network has developed an evidence-based practice which may be suitable for use in a BH SBIRT program. The Trauma Adaptive Recovery Group Education and Therapy for Adolescents and Pre-Adolescents (TARGET-A) has been evaluated in 248 clinical trials with control groups and can be completed in as little as 4 sessions. This intervention is designed for groups and/or individual children, adolescents and their parents that is easily adapted to settings where youth or families enter and leave services rapidly (NCTSN, 2008).

The prevalence of issues such as depression, anxiety, and trauma among primary care patients call for further exploration to determine if certain SBIRT components may be applied to symptoms of these disorders among medical patients. These findings also highlight the value of universal screening, a principal component of SBIRT, in addressing mental health issues in primary care and other health care settings.

VIII. IMPLICATIONS FOR FUTURE PROGRAMS

While there is substantial research for the effectiveness of SBIRT in reducing unhealthy alcohol use and tobacco use/misuse, the evidence for similar models in addressing drug abuse and mental health conditions such as depression, anxiety and trauma is still being developed. As such, SAMHSA would recommend investment in developing SBIRT-like models for most common behavioral health conditions, for use in public health settings. This would involve services research, demonstrations, and conducting rigorous comparative effectiveness evaluations of behavioral health SBIRT programs beyond those already proven effective for alcohol or tobacco, in possible collaboration with NIMH, NIAAA and/or NIDA.

Numerous screening and intervention programs in a variety of settings and populations have recently defined themselves as “SBIRT programs.” Most often these programs do not meet the criteria established in this paper to be designated as a comprehensive SBIRT model. Both a strong research base and more consistent terminology and definitions for what constitutes a true SBIRT model are lacking. Although SBIRT and its components have been utilized across programs, the effectiveness of SBIRT programs can vary in their fidelity, application, and comprehensiveness.

In considering the future of SBIRT program implementation, some or all of the following could be pursued:

- Partnership with one or more external, authoritative bodies. This may involve approaching the US Preventative Services Task Force to develop an SBIRT definition and/or taxonomy which reflects the latest science-based approach and is vetted with the field.
- Collaboration with NIH (NIDA, NIMH) and/or AHRQ to conduct more research on SBIRT approaches for drug abuse, depression, anxiety, trauma, etc., to help establish parameters that are critical to effective implementation.
- Diversifying the SAMHSA SBIRT program portfolio and dedicating increased evaluation resources to examine the value of complementing SBIRT for alcohol and drugs with screening and intervention for other behavioral health conditions.

References

- Amaral, M.B., Ronzani, T.M., Souza-Formigoni, Maria, L.O. (2010). Process evaluation of the implementation of a screening and brief intervention program for alcohol risk in primary health care: An experience in Brazil. *Drug and Alcohol Review*, 29 (2), 162-168.
- Amaro, H., Reed, E., Rowe, E., Picci, J., Mantella, P., et al. (2010). Brief screening and intervention for alcohol and drug use in a college student health clinic: Feasibility, implementation, and outcomes. *Journal of American College Health*, 58(4), 357-364.
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders* (4th ed., text revision). Washington, DC: Author.
- APA Achievement Awards (2010). A new direction in depression treatment in Minnesota, DIAMOND program, Institute for Clinical Systems Improvement, Bloomington, Minnesota. *Psychiatric Services*, 61(10), 1042-1044.
- Arndt, S., Schultz, S. K., Turvey, C., & Petersen, A. (2002). Screening for alcoholism in the primary care setting: Are we talking to the right people? *Journal of Family Practice*, 51(1), 41-46.
- Babor, T. F., McRee, B. G., Kassebaum, P. A., Grimaldi, P. L., Ahmed, K., & Bray, J. (2007). Screening, Brief Intervention, and Referral to Treatment (SBIRT): Toward a public health approach to the management of substance abuse. *Substance Abuse*, 28(3), 7-30.
- Babor, T.F. (2008). Taking stock: Twenty-five years of translational research on alcohol screening and brief intervention. *NAT: Nordisk Alkohol & Narkotikatidskrift*, 25(6), 578-580.
- Babor, T. F., & Higgins-Biddle, J. C. (2000). Alcohol screening and brief intervention: Dissemination strategies for medical practice and public health. *Addiction*, 95, 677-686.
- Babor, T. F., Higgins-Biddle J., Dauser, D., Higgins P, & Bureson, J. (2005). Alcohol screening and brief intervention in primary care settings: Implementation models and predictors. *Journal on Studies of Alcohol and Drugs*, 66(3), 361-369.
- Babor, T., Higgins-Biddle, J., Higgins, P., Gassman, R., Gould, B. Training Medical Providers to Conduct Alcohol Screening and Brief Interventions. *Substance Abuse*, 25 (1), 17-26.
- Babor, T., McRee, B., Kassebaum, P., Grimaldi, P., Ahmed, K., Bray, J. (2007). Screening, Brief Intervention, and Referral to Treatment (SBIRT). *Substance Abuse*, 28 (3), 7-30.
- Baer, J.S., Kivlahan, D.R., Blume, A.W., McKnight, P., & Marlatt, A. (2001). Brief intervention for heavy-drinking college students: 4 Year follow-up and natural history. *American Journal of Public Health*, 91(8), 1310-1316.

- Baker, A., Lee, N. K., Claire, M., Lewin, T. J., Grant, T., & Pohlman, S. (2005). Brief cognitive behavioural interventions for regular amphetamine users: A step in the right direction. *Addiction*, *100*, 367–378.
- Barry, D. (2002). Brief Alcohol Screening and Intervention for College Students (BASICS): A Harm Reduction Approach. *Journal of Psychiatry & Law*, *30* (2), 275-9.
- Begun, A. L., Rose, S.J., LeBel, T. P., & Teske-Young, B. A. (2009). Implementing Substance Abuse Screening and Brief Motivational Intervention With Women in Jail. *Journal of Social Work Practices in the Addictions*, *9*(1), 113–131.
- Beich, A., Gannik, D., Malterud, K. (2002). Screening and brief intervention for excessive alcohol use: Qualitative interview study of the experiences of general practitioners. *British Medical Journal*, *325* (7369), 870-870.
- Beich, A., Thorsen, T., & Rollnick, S. (2003). Screening in brief intervention trials targeting excessive drinkers in general practice: Systematic review and meta-analysis. *BMJ*, *327*, 536–542.
- Bernstein, E., Bernstein, J., & Levenson, S. (1997). Project ASSERT: An ED-based intervention to increase access to primary care, preventive services, and the substance abuse treatment system. *Annals of Emergency Medicine*, *30*, 181–189.
- Bernstein, J., Bernstein, E., Tassiopoulos, K., Heeren, T., Levenson, S., & Hingson, R. (2005). Brief motivational intervention at a clinic visit reduces cocaine and heroin use. *Drug Alcohol Dependence*, *77*(1), 49–59.
- Bernstein E., Topp D., Shaw E., Girard C., Pressman K., Woolcock E., Bernstein J., (2009). A preliminary report of knowledge translation: Lessons from taking screening and brief intervention techniques from the research setting into regional systems of care. *Academic Emergency Medicine*, *16* (11), 1225-33.
- Bertholet, N., Daeppen, J.-B., Wietlisbach, V., Fleming, M., & Burnand, B. (2005). Reduction of alcohol consumption by brief alcohol intervention in primary care: systematic review and meta-analysis. *Archives of Internal Medicine* *165*, 986–995.
- Bien, T., Miller, W. R., & Tonigan, J. S. (1993). Brief interventions for alcohol problems: A review. *Addiction*, *88*, 315–336.
- Blume, A.W.; Marlatt, G. A. (2004). Motivational Enhancement as a Brief Intervention for College Student Drinkers. In: *Handbook of Motivational Counseling: Concepts, Approaches, and Assessment*, 409-420. New York, NY: John Wiley & Sons Ltd.
- Bondy, S. J., Rehm, J., Ashley, M. J., Walsh, G., Single, E., & Room, R. (1990). Low-risk drinking guidelines: the scientific evidence. *Canadian Journal of Public Health*, *90*(4), 264–70. [PMID: 10489725]

- Borsari, B.; Carey, K. B. (2000). Effects of a brief motivational intervention with college students. *Journal of Consulting and Clinical Psychology*, 68(4),728-733.
- Bradley, K. A., DeBenedetti, A. F., Volk, R. J., Williams, E. C., Frank, D., & Kivlahan, D. R. (2007). AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcoholism, Clinical and Experimental Research*, 31, 1208–1217.
- Bray, J., Mills, M., Bray, Lessell M., Lennox, R., McRee, B., Goehner, D., Higgins-Biddle, J. (2009). Evaluating web-based training for employee assistance program counselors on the use of screening and brief intervention for at-risk alcohol use. *Journal of Workplace Behavioral Health*, 24 (3). 307-319.
- Bray, J.W., Zarkin, G.A., Davis, K.L., Mitra ,D., Higgins-Biddle, J.C., Babor, T.F. (2007). The effect of screening and brief intervention for risky drinking on health care utilization in managed care organizations. (eng), *Medical Care*, 45 (2), 177-82.
- Brewin, C. (2005). Systematic review of screening instruments for adults at risk of PTSD. *Journal of Traumatic Stress*. Feb; 18(1): 53-62
- Briere, J. (1995). Trauma Symptom Inventory professional manual. Odessa, FL: Psychological Assessment Resources
- Brown, R.L., Fleming, M.F., (1998). Training the trainers: substance abuse screening and intervention. (eng), *International Journal of Psychiatry in Medicine*, 28 (1), 137-46.
- Brown, R. L., Saunders, L. A., Bobula, J. A., Mundt, M. P., & Koch, P. E. (2007). Randomized-controlled trial of a telephone and mail intervention for alcohol use disorders: Three-month drinking outcomes. *Alcoholism, Clinical and Experimental Research*, 31(8), 1372–1379.
- Bruce, M. L., Ten Have, T. R., Reynolds III C.F., et al. (2004). Reducing suicidal ideation and depressive symptoms in depressed older primary care patients: a randomized controlled trial. *JAMA*, 294, 1081-1091.
- Cameron, Jacqui; Lee, Nicole; Harney, Angela. (2010). Changes in attitude to, and confidence in, working with comorbidity after training in screening and brief intervention. *Mental Health and Substance Use: Dual Diagnosis*, 3(2), 24-130.
- Casset J.C., Bérenger P., Bosson J.L., Lacroix A., (2008). Evaluation à 1 an de l'intervention brève pratiquée par des médecins généralistes chez des patients ayant un mésusage d'alcool. *La Revue Du Praticien*, 58 (12), 25-31.
- Center for Substance Abuse Treatment. (1999). *Brief interventions and brief therapies for substance abuse*. Treatment Improvement Protocol 34. HHS Publication No. (SMA) 99-3353. Rockville, MD: Substance Abuse and Mental Health Services Administration.

- Charbonney, Emmanuel (2009). Alcohol, drugs and trauma: Consequences, screening and intervention in 2009. *Trauma*, 12 (1) 5-12.
- Chisholm, D. J., Young, R. R., & McAlearney, A. S. (2005). Implementation of a touch-screen new patient registration system: A case study. *Journal of Medical Practice & Management*, 21(3), 159–162.
- Christensen, M.H., Boisse, N., Sanchez, W., Friedmann, P.D. (2004). Enhancing vocational rehabilitation counselors' substance abuse screening and brief intervention practices. *Journal of Vocational Rehabilitation*, 21 (3), 157-163.
- Cole, M. G., Dendukuri, N. (2004). The feasibility and effectiveness of brief interventions to prevent depression in older subjects: a systematic review. *International Journal of Geriatric Psychiatry*, 19, 1019-1025.
- Copeland, J., Swift, W., Roffman, R., & Stephens, R. (2001.) A randomized controlled trial of brief cognitive-behavioral interventions for cannabis use disorder. *Journal of Substance Abuse Treatment*. 21(2), 55–64.
- Corrao, G., Bagnardi, V., Zambon, A., & Arico, S. (1999). Exploring the dose-response relationship between alcohol consumption and the risk of several alcohol-related conditions: a meta-analysis. *Addiction*, 94, 1551–1573.
- Covell, N., Essock, S., Felton, C., and Donahue, S. (2006). Characteristics of Project Liberty clients that predicted referrals to intensive mental health services. *Psychiatric Services*, 57(9), 1313-1315.
- Crawford et al. (2010). The effect of referral for brief intervention for alcohol misuse on repetition of deliberate self-harm: an exploratory randomized controlled trial. *Psychological Medicine*, 40, 1-8.
- [Cucciare MA, Weingardt KR, Humphreys K.](#) (2009). How Internet technology can improve the quality of care for substance use disorders. *Curr Drug Abuse Rev*, Sep;2(3):256-2.
- Cuijpers, P., van Straten, A., van Schaik, A., Andersson, G., (2009). Psychological treatment of depression in primary care: a meta-analysis. *British Journal of General Practice*, 59(559), 51-60.
- Cunningham, R., Bernstein, S., Walton, M., Broderick, K., Vaca, F., & Woolard, R., et al. (2009). Alcohol, Tobacco, and Other Drugs: Future Directions for Screening and Intervention in the Emergency Department. *Academic Emergency Medicine*, 16:1078–1088
- Daepfen, J. B., Gaume, J., Bady, P., Yersin, B., Calmes, J. M., & Givel, J. C., et al. (2007). Brief alcohol intervention and alcohol assessment do not influence alcohol use in injured patients treated in the emergency department: A randomized controlled clinical trial. *Addiction*, 102, 1224 –1233.

- Donker, T., Griffiths, K.M., Cuijpers, P., & Christensen, H. (2009) Psychoeducation for depression, anxiety and psychological distress: a meta-analysis. *BMC Medicine*, 7(79).
- De Micheli, D, Fisberg M, Formigoni ML. (2004). Study on the effectiveness of brief intervention for alcohol and other drug use directed to adolescents in a primary health care unit. *Rev Assoc Med Bras*, 50:305-13.
- Derogatis, L. R. (2001). Brief Symptom Inventory (BSI)-18: Administration, scoring and procedures manual. Minneapolis, MN: NCS Pearson. D'Onofrio, G., & Degutis, L. C. (2002). Preventive care in the emergency department: Screening and brief intervention for alcohol problems in the emergency department: A systematic review. *Academic Emergency Medicine*, 9, 627–638.
- D'Onofrio, G., Bernstein, E., Bernstein, J., Woolard, R. H., Brewer, P. A., & Craig, S. A., et al. (1998). Patients with alcohol problems in the emergency department, part 2: Intervention and referral. SAEM Substance Abuse Task Force. *Academic Emergency Medicine*, 5, 1210–1217.
- Diehl, A., Nakovics, H., Bernhard, C., Reinhard, I., Kiefer, F., & Mann, K. (2007). Consultation-liaison psychiatry in general hospitals: improvement in physicians' detection rates of alcohol use disorders. *Journal of Psychosomatics*, 50, 599-604.
- Donker, T., Griffiths, K., Cuijpers, P., Christensen, H. (2009). Psychoeducation for depression, anxiety and psychological distress: A meta-analysis. *BMC Medicine* 7(79).
- Dube, SR, Williamson, DF, Thompson, T, Felitti, VJ, Anda, RF (2004). Assessing the reliability of retrospective reports of adverse childhood experiences among adult HMO members attending a primary care clinic. *Child Abuse and Neglect*, 28, 729-737.
- Dimidjian, S., et al. (2006). "Randomized Trial of Behavioral Activation, Cognitive Therapy, and Antidepressant Medication in the Acute Treatment of Adults With Major Depression". [*Journal of Consulting and Clinical Psychology* 74 \(4\): 658–670.](#)
- Dunn, C. W., & Ries, R. (1997). Linking substance abuse services with general medical care: Integrated, brief interventions with hospitalized patients. *American Journal of Drug and Alcohol Abuse*, 23, 1–13.
- Dyches, H., Alemagno, S., Llorens, S. A., & Butts, J. M. (1999). Automated telephone-administered substance abuse screening for adults in primary care. *Health Care Management Science*, 2(4), 199–204.
- Elvy, G. A., Wells, J. E., & Baird, K. A. (1988). Attempted referral as intervention for problem drinking in the general hospital. *British Journal of Addiction*, 83, 83–89.

- Emmen, M. J., Schippers, G. M., Bleijenberg, G., & Wollersheim, H. (2004). Effectiveness of opportunistic brief interventions for problem drinking in a general hospital setting: Systematic review. *BMJ*, *328*(7435), 318.
- Falk, D. E., Yi, H. Y., & Hiller-Sturmhöfel, S. (2006). An epidemiologic analysis of co-occurring alcohol and tobacco use and disorders: Findings from the National Epidemiologic Survey on Alcohol and Related Conditions. *Alcohol Research & Health*, *29*, 162–171.
- Ford, JD, and Russo, E. (2006). Trauma-focused, present-centered, emotional self-regulation approach to integrated treatment for posttraumatic stress and addiction: Trauma adaptive recover group education and therapy (TARGET). *American Journal of Psychotherapy*, *60*(4), 335-355.
- Fleming, M. F., Barry, K. L., Manwell, L. B., Johnson, K., & London, R. (1997). Brief physician advice for problem alcohol drinkers: A randomized controlled trial in community-based primary care practices. *JAMA*, *277*, 1039–1045.
- Fleming M.F., Graham A.W. (2001). Screening and brief interventions for alcohol use disorders in managed care settings. (eng), *Recent Developments in Alcoholism: An Official Publication of The American Medical Society on Alcoholism, The Research Society on Alcoholism, and The National Council on Alcoholism*, *15*, 393-416.
- Fleming, M.F.; Mundt, M.P.; French, M.T., Manwell, L. B., Stauffacher, E. A., Barry, K. L. (2002). Brief physician advice for problem drinkers: Long-term efficacy and benefit-cost analysis. *Alcoholism: Clinical and Experimental Research*, *26*(1), 36–43.
- Fornili, K., Alemi, F. (2007). Medicaid Reimbursement for Screening and Brief Intervention: Amending the Medicaid State Plan and Approving State Appropriations for the Medicaid State Match. *Journal of Addictions Nursing*, *18* (4), 225-232.
- Foy, J., ed. (2010). Enhancing pediatric mental health care: Report from the American Academy of Pediatrics task force on mental health. *Pediatrics*, June, *125*, supplement 3, S69-S160.
- Funderburk, J. S., Maisto, S. A., & Sugarman, D. (2008). Brief alcohol intervention and multiple risk factors in primary care. *Substance Abuse*, *28*(4), 93–105.
- Furtado, E. F.; Corradi-Webster, C. M.; Laprega, M. R. (2008). Implementing brief interventions for alcohol problems in the public health system in the region of Ribeirao Preto, Brazil: Evaluation of the PAI-PAD training model. *NAT: Nordisk Alkohol & Narkotikatidskrift*, *25*(6), 539-551.
- Gallo, J., Zubritsky, C., Maxwell, J., et al. (2004). Primary care clinicians evaluate integrated and referral models of behavioral health care for older adults: results from a multisite effectiveness trial (PRISM-E). *Annals of Family Medicine*, *2*(4).

- Gaynes et al. (2010). Feasibility and diagnostic validity of the M-3 checklist: a brief, self-rated screen for depressive, bipolar, anxiety, and post-traumatic stress disorders in primary care. *Annals of Family Medicine*, Mar-Apr, 8(2): 160-9.
- Gentilello, L. M., Donovan, D. M., Dunn, C. W., Rivara, F. P. (1999). Alcohol interventions in a trauma center as a means of reducing the risk of injury recurrence. *Annals of Surgery*, 230, 1-18.
- Gilbody, S., House, A., Sheldon, T., (2009). Screening and case finding instruments for depression (review). *The Cochrane Library*, 3, 1-30.
- Grupp-Phelan, J., et al. (2009). Referral and resource use patterns for psychiatric-related visits to pediatric emergency departments. *Pediatric Emergency Care*, 25(4), 217-220.
- Guevara, J., Greenbaum, P., Shera, D., Bauer, L., and Scharz D. (2009). Survey of mental health consultation and referral among primary care physicians. (2009). *Academic Pediatrics*, 9(2), 123-127.
- Hanewinkel, Reiner; Wiborg, Gudrun (2005). Brief alcohol screening and intervention for college students (BASICS): A German pilot study. *Sucht: Zeitschrift für Wissenschaft und Praxis*, 51(5), 285-290.
- Hansen, M., Andersen, T. Armour, C Elklit, A., Palic, S., and Mackrill, T. (2010). PTSD-8: A short PTSD inventory. *Clinical Practice & Epidemiology in Mental Health*, 6, 101-108
- Heather, Nick; Dallolio, Emma; Hutchings, Deborah; Kaner, Eileen; White, Martin (2004). Implementing routine screening and brief alcohol intervention in primary health care: A Delphi survey of expert opinion. *Journal of Substance Abuse*, 9(2), 68-85.
- Henslee, A.M. (2009) Providing personalized feedback regarding alcohol use in a group format to college freshmen. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 69(10-B), 6414.
- Hillman, A., McCann, B., & Walker, N. P. (2001). Specialist alcohol liaison services in general hospitals improve engagement in alcohol rehabilitation and treatment outcome. *Health Bulletin*, 59, 420-423.
- Hungerford, D.W., Williams, J.M., Furbee, P.M., Manley, W.G. 3rd, Helmkamp, J.C., Horn, K., Pollock, D.A. (2003). Feasibility of screening and intervention for alcohol problems among young adults in the ED. *The American Journal of Emergency Medicine*, 21 (1) 14-22.
- Hungerford, D. W., & Pollock, D. A. (2003). Emergency department services for patients with alcohol problems: Research directions. *Academy of Emergency Medicine*, 10(1), 79-84.
- Kahan, M., Wilson, L., & Becker, L. (1995). Effectiveness of physician-based interventions with problem drinkers: A review. *Canadian Medical Association Journal*, 152, 851-859.

- Kaner, E. F. S., Beyer, F., Dickinson, H. O., Pienaar, E., Campbell, F., Schlesinger, C., et al. (2009). Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database of Systematic Review*, 18(2), CD004148.
- Kaner, E. F., Dickinson, H. O., Beyer, F., Pienaar, E., Schlesinger, C., & Campbell, F., et al. (2009). The effectiveness of brief alcohol interventions in primary care settings: A systematic review. *Drug and Alcohol Review*, 28(3), 301–323.
- Kroenke K, Spitzer R L, Williams J B. The PHQ-9: validity of a brief depression severity measure. *Journal of General Internal Medicine* 2001; 16(9): 606-613
- Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: Validity of a two-item depression screener. *Med Care*. 2003; 41(11):1284-1292.
- Kraemer, K. (2007). The Cost-Effectiveness and Cost-Benefit of Screening and Brief Intervention for Unhealthy Alcohol Use in Medical Settings. *Substance Abuse*, 28 (3), 67-77.
- Kunz F.M. Jr, French M.T., Bazargan-Hejazi S. (2004). Cost-effectiveness analysis of a brief intervention delivered to problem drinkers presenting at an inner-city hospital emergency department. *Journal of Studies on Alcohol*, 65 (3), 363-70.
- Larimer, Mary E.; Cronce, Jessica M.; Lee, Christine M.; Kilmer, Jason R. (2004). Brief Intervention in College Settings. *Alcohol Research & Health*, 28 (2), 94-104.
- Levkoff, S. E., Chen, H., Coakley E., et al. (2002). Design and sample characteristics of the PRISM-E multisite randomized trail to improve behavioral health care for the elderly. *Journal of Aging Health*, 16, 3-27.
- Levy, S., & Knight, J. R. (2008). Screening, brief intervention, and referral to treatment for adolescents. *Journal of Addiction Medicine*, 2(4), 215–221
- Longabaugh, R., Woolard, R. F., Nirenberg, T. D., Minugh, A. P., Becker, B., & Clifford, P. R., et al. (2001). Evaluating the effects of a brief motivational intervention for injured drinkers in the emergency department. *Journal on Studies of Alcohol and Drugs*, 62, 806–16.
- Ludman, E., Katon, W., Bush T., Rutter, C., Lin, E., & Simon, G., et al. (2003). Behavioral factors associated with symptom outcomes in a primary care-based depression prevention intervention trial. *Psychological Medicine*, 33, 1061-1070.
- Luoma, J, Martin, C., Pearson, J. (2002) Contact with mental health and primary care providers before suicide: a review of the evidence. *American Journal of Psychiatry*. Jun,159(6) :909-16.
- Madras, B. K., Compton, W. M., Avula, D., Stegbauer, T., Stein, J. B., & Clark, H. W. (2008). Screening, brief interventions, referral to treatment (SBIRT) for illicit drug and alcohol use at multiple healthcare sites: Comparison at intake and 6 months later. *Drug Alcohol Dependence*, doi:10.1016/j.drugalcdep.2008.08.003

- Magill M, Barnett NP, Apodaca TR, Rohsenow DJ, Monti PM. (2009). The role of marijuana use in brief motivational intervention with young adult drinkers treated in an emergency department. *J Stud Alcohol Drugs*.70 (3):409-13.
- Mallett, K., Bachrach, R., Turrisi, R. (2009). Examining the Unique Influence of Interpersonal and Intrapersonal Drinking Perceptions on Alcohol Consumption Among College Students. *Journal of Studies on Alcohol and Drugs*, 70 (2), 178-185.
- Marlatt, G. A., Baer, J. S., Dimeff, L. A., Larimer, M. E., Quigley, L. A., & Somers, J. M., et al. (1998). Screening and brief intervention for high-risk college student drinkers: Results from a 2-year follow-up assessment. *Journal of Consulting and Clinical Psychology*, 66, 604-615.
- Marsden, J., Stillwell, G., Barlow, H., Boys, A., Taylor, C., & Hunt, N., et al. (2006). An evaluation of a brief motivational intervention among young ecstasy and cocaine users: no effect on substance and alcohol use outcomes. *Addiction*. 101(7), 1014–1026.
- Martens, M. P.; Cimini, M. D.; Barr, A. R.; Rivero, E. M.; Vellis, P.A.; Desemone, G. A.; Horner, K. J. (2007). Implementing a screening and brief intervention for high-risk drinking in university-based health and mental health care settings: Reductions in alcohol use and correlates of success. *Addictive Behaviors* 32(11), 2563-2572.
- Mastroleo, Nadine R. (2009). Comparison of supervision training techniques in a motivational enhancement intervention on college student drinking. *Dissertation Abstracts International Section A: Humanities and Social Sciences*, 69(11-A), 4254.
- McNaughton, J. (2009). Brief interventions for depression in primary care: a systematic review. *Canadian Family Physician*, Aug , 55(8) 789-96.
- McPherson, T. L, Goplerud, E., Olufokunbi-Sam, D., Jacobus-Kantor, L., Lusby-Treber, K. A., et al. (2009). Workplace Alcohol Screening, Brief Intervention, and Referral to Treatment (SBIRT): A Survey of Employer and Vendor Practices. *Journal of Workplace Behavioral Health*, 4(3), 285–306.
- McQueen, J., Howe, T. E., Allan, L., & Mains, D. (2009). Brief interventions for heavy alcohol users admitted to general hospital wards (review). *Cochrane Database Syst Rev*, 3:CD005191.
- McRee, B., Babor, T.F., Maranjian Church, O. (2002). Instructor's Manual for Alcohol Screening & Brief Intervention. *Instructor's Manual for Alcohol Screening & Brief Intervention*, 61.
- McRee, B., Granger, J., Babor T. F., Feder, I., Folino, M., et al. (2005). Reducing tobacco use and risky drinking in underserved populations: The need for better implementation models. *Annals of Family Medicine*, 3(2), 558–560

- Mertens, J. R., Weisner, C., Ray, G. T., Fireman, B., & Walsh, K. (2005). Hazardous drinkers and drug users in HMO primary care: prevalence, medical conditions, and costs. *Alcoholism, Clinical and Experimental Research*, 29, 989–998.
- Meyer, F., Peteet, J., Joseph R. (2009). Models for co-occurring mental and medical disorders. *Harvard Review of Psychiatry*, Nov/Dec, 353-360.
- Miller, W. R., & Rollnick, S. (2002). *Motivational interviewing: Preparing people for change*. (2nd ed.) New York, NY: The Guilford Press.
- Modesto-Lowe, V., & Boornazian, A. (2000). Screening and brief intervention in the management of early problem drinkers: Integration into healthcare settings. *Disease Management & Health Outcomes*, 8(3), 129–137.
- Moos, R. H., & Moos, B. S. (2003). Long-term influence of duration and intensity of treatment on previously untreated individuals with alcohol use disorders. *Addiction*, 98(3), 325–337.
- Morrissey, J., Jackson, E., Ellis, A., Amaro, H., Brown, V., & Najavits L. (2005). Twelve-month outcomes of trauma-informed interventions for women with co-occurring disorders. *Psychiatric Services*, 56(10), 1213-1221.
- Moyer, A., Finney, J., Swearingen, C., & Vergun, P. (2002). Brief interventions for alcohol problems: A meta-analytic review of controlled investigations in treatment-seeking and non-treatment-seeking populations. *Addiction*, 97, 279–292.
- Murphy, J. G., Duchnick, J. J., Vuchinich, R. E., Davison, J. W., Karg, R. S. et al. (2001). Relative efficacy of a brief motivational intervention for college student drinkers. *Psychology of Addictive Behaviors*, 15(4), 373–379.
- National Institute on Alcohol Abuse and Alcoholism (NIAAA). (2005a). *Alcohol alert number 66: Brief interventions*. Rockville, MD: U.S. Department of Health and Human Services, National Institute on Alcohol Abuse and Alcoholism. Retrieved June 4, 2010 from <http://pubs.niaaa.nih.gov/publications/AA66/AA66.pdf>.
- Nisenson, LG, Pepper, CM, Schwenk, TL & Coyne, JC (1998). The nature and prevalence of anxiety disorders in primary care. *General Hospital Psychiatry*, 20, 21-28.
- O'Brien, M. C., McCoy, T. P., Champion, H., Mitra, A., Robbins, A., & Teuschlser, et al. (2006). Single question about drunkenness to detect college students at risk for injury. *Academic Emergency Medicine*, 13(6), 629–636
- Olfson, M., Shea, S., Feder, A., Fuentes, M., Nomura, Y., & Gameroff, M., et al. (2000). Prevalence of anxiety, depression, and substance use disorders in an urban general medicine practice. *Archives of Family Medicine*, 9(9), 876–883.

- Osilla, K.C.; Cruz, E. dela; Miles, J. N.V.; Zellmer, S.; Watkins, K.; Larimer, M. E.; Marlatt, G. A.. (2010). Exploring productivity outcomes from a brief intervention for at-risk drinking in an employee assistance program. *Addictive Behaviors*, 35 (3) 194-200.
- Oslin, D. W., Sayers, S., Ross, J., et al. (2003). Disease management for depression and at-risk drinking via telephone in an older population of veterans. *Psychosomatic Medicine*, 65, 931-937.
- Palmer, et al. (2003). Screening for depression in medical care: Pitfalls, alternatives, and revised priorities. *Journal of Psychosomatic Research*, 54, 279-287.
- Paperny, D. M. N. (1997). Computerized health assessment and education for adolescent HIV and STD prevention in health care settings and schools. *Health Education & Behavior*, 24(1), 54-70.
- Paperny, D. M. N., Aono, J., Lehman, R., Hammar, S., & Risser, J. (1990). Computer-assisted detection and intervention in adolescent high-risk health behaviors. *Journal of Pediatrics*, 116(3), 453-462.
- Peltzer K., Seoka, P., Babor, T., Obot, I. (2006). Training primary care nurses to conduct alcohol screening and brief interventions in South Africa. (eng), *Curationis*, 29 (2), 16-21.
- Prevention of "Risky" Drinking among Students at a Brazilian University Alcohol and Alcoholism: International Journal of the Medical Council on Alcoholism. *International Journal of the Medical Council on Alcoholism*, (2008). 43 (4), 470-476.
- Prins A, Ouimette P, Kimerling R et al. (2003). The primary care PTSD screen (PC-PTSD): Development and operating characteristics. *Primary Care Psychiatry*, 9: 9-14.
- Regar, M. A., Gahm, G. A., Swanson, R. D., Duma, S. J. (2009). Association between number of deployments to Iraq and mental health screening outcomes in US Army soldiers. *Journal of Clinical Psychiatry*, 70(9), 1266-1272.
- Reid, M. C., Fiellin, D. A., & O'Connor, P. G. (1999). Hazardous and harmful alcohol consumption in primary care. *Archives of Internal Medicine*, 159(15), 1681-1689. [PMID: 10448769]
- Resick, PA (2001). Cognitive therapy for posttraumatic stress disorder. *Journal of Cognitive Psychotherapy: An International Quarterly*, 15(4), 321-329.
- Richmond, R., Heather, N., Wodak, A., Kehoe, L., & Webster, I. (1995). Controlled evaluation of a general practice-based brief intervention for excessive drinking. *Addiction*, 90(1), 119-132.
- Rinfrette, E. S. (2009). Treatment of anxiety, depression, and alcohol disorders in the elderly: social work collaboration in primary care. *Journal of Evidence-Based Social Work*, 6, 79-91.

- Rivara, F. P., Tollefson, S., Tesh, E., & Gentilello, L. M. (2000). Screening trauma patients for alcohol problems: Are insurance companies barriers? *Journal of Trauma*, 48(1), 115–118.
- Roberts, N. P., Kitchiner, N. J., Kenardy, J., & Bisson, J. I. (2009). Systematic review and meta-analysis of multiple-session early interventions following traumatic events. *American Journal of Psychiatry*, 166 (3), 293-301
- Roche, A. M., & Freeman, T. (2004). Brief interventions: Good in theory but weak in practice. *Drug and Alcohol Review*, 23(1), 11–18.
- Ronzani, T. M., do Amaral, M., Bitarello, Souza-Formigoni, M. L. O., Babon, T. F. (2008). Evaluation of a training program to implement alcohol screening, brief intervention and referral to treatment in primary health care in Minas Gerais, Brazil. *NAT: Nordisk Alcohol & Narkotikatidskrift*, 25(6), 529-538.
- Ronzani T.M., Mota D.C., Souza I.C., (2009). Alcohol prevention within primary care in municipalities in the state of Minas Gerais, Southeastern Brazil. *Revista De Saúde Pública*, (43), 51-61.
- Roy-Byrne, P., Craske, M. G., Sullivan, G., Rose, R. D., Edlund, M. J., & Lang, A. J. et al. (2010). Delivery of evidence-based treatment for multiple anxiety disorders in primary care a randomized controlled trial. *JAMA*, 303(19), 1921–1928.
- Rueda S, Park-Wyllie LY, Bayoumi AM, Tynan AM, Antoniou TA, & Rourke SB, et al. (2006). Patient support and education for promoting adherence to highly active antiretroviral therapy for HIV/AIDS. *Cochrane Database Syst Rev*. 19 (3):CD001442.
- Saitz, R. (2010). Candidate Performance Measures for Screening for, Assessing, and Treating Unhealthy Substance Use in Hospitals: Advocacy or Evidence-Based Practice? Commentary from the *Ann Intern Med*. 2010;153:40-43.
- Saitz, R., Helmuth, E. D., Aromaa, S. E., Guard, A., Belanger, M., et al. (2004). Web-based screening and brief intervention for the spectrum of alcohol problems. *Preventive Medicine*, 39, 969–975.
- Saitz, R., Horton, N. J., Sullivan, L. M., Moskowitz, M. A., & Samet, J. H. (2003). Addressing alcohol problems in primary care: A cluster randomized, controlled trial of a systems intervention. *Annals of Internal Medicine*, 138(5), 372–382.
- Saitz R, Svikis D, D’Onofrio G, et al. Challenges applying alcohol brief intervention in diverse practice settings: Populations, outcomes, and costs. *Alcohol Clin Exp Res*. 2006; 30:332–338.
- Saitz, R., Palfi, T. P., Cheng, D. M., Horton, N. J., Freedner, N., & Dukes, K. et al. (2007). Brief intervention for medical inpatients with unhealthy alcohol use: a randomized, controlled trial. *Annals of Internal Medicine*, 146, 167–176.

- SAMHSA's National Registry for Evidence-Based Programs and Practices. (2008). Brief alcohol screening and intervention for college students (basics). Retrieved on June 4, 2010 from http://nrepp.samhsa.gov/programfulldetails.asp?PROGRAM_ID=156
- Schermer C.R. (2005). Feasibility of alcohol screening and brief intervention. (eng). *The Journal of Trauma*, 59, S119-23.
- Schermer, C. R., Gentilello, L. M., & Hoyt, D. B. (2003). National survey of trauma surgeons' use of alcohol screening and brief intervention. *Journal of Trauma*, 55, 849–855.
- Schonfeld L, King-Kallimanis B.L., Duchene D.M., Etheridge R.L., Herrera J.R., Barry K.L., Lynn N. (2010). Screening and brief intervention for substance misuse among older adults: The Florida BRITE project. (eng), *American Journal of Public Health*, 100 (1), 108-14.
- Schwartz. Allan J. (2006). College student suicide in the United States: 1990-1991 through 2003-2004. *Journal of American College Health*, 54 (6), 341-352.
- Seale, J.P. Monteiro, M.G. (2008). The dissemination of screening and brief intervention for alcohol problems in developing countries: Lessons from Brazil and South Africa. *NAT: Nordisk Alkohol & Narkotikatidskrift*, 25(6), 565-577.
- Seale J.P., Shellenberger S., Boltri J.M., Okosun I.S., Barton B. (2005). Effects of screening and brief intervention training on resident and faculty alcohol intervention behaviours: A pre-post-intervention assessment. *BCM Family Practice*, 6, 46.
- Shelbourne, CD, Jackson, CA, Meredith, LS, Camp P, & Wells KB (1996). Prevalence of comorbid anxiety disorders in primary care outpatients. *Archives of Family Medicine*, 5(1), 27-34.
- Shellenberger S., Seale J.P., Harris D.L., Johnson J.A., Dodrill C.L., Velasquez M.M., (2009) Applying team-based learning in primary care residency programs to increase patient alcohol screenings and brief interventions. *Journal of the Association of American Medical Colleges*, 84 (3), 340-6.
- Simon, G. E., Von Korff, M., Ludeman, E. J., et al. (2002). Cost-effectiveness of a program to prevent depression relapse in primary care. *Med Care*, 40, 941-950.
- Simon, G. E., Katon, W. J., VonKorff, et al. (2001). Cost-effectiveness of a collaborative care program for primary care patients with persistent depression. *American Journal of Psychiatry*, 158, 1638-1644.
- Smith, P.C., Schmidt, S.M., Allensworth-Davies, D., & Saitz, R. (2010) A single-question screening test for drug use in primary care. *Archives of Internal Medicine*, 170(13):1155-1160.

- Souza-Formigoni, M. L. O.; Boemgen-Lacerda, R.; Vianna, V. P. T. (2008). Implementation of alcohol Screening and brief intervention in primary care units in two Brazilian states: A case study. *NAT: Nordisk Alkohol & Narkotikatidskrift*, 25(6), 533-564.
- Spandorfer, J. M., Israel, Y., & Turner, B. J. (1999). Primary care physicians' views on screening and management of alcohol abuse: Inconsistencies with national guidelines. *Journal of Family Practice*, 48(11), 899-902.
- Stephens R. S., Roffman R. A., & Curtin L. (2000). Comparison of extended versus brief treatments for marijuana use. *Journal of Consulting and Clinical Psychology*, 68, 898-908.
- Tait, R. J., Hulse, G. K., & Robertson, S. I. (2004). Effectiveness of a brief intervention and continuity of care in enhancing attendance for treatment by adolescent substance users. *Drug Alcohol Dependence*, 74(3), 289-296.
- Tait, R. J., Hulse, G. K., Robertson, S. I., & Sprivulis, P. C. (2005). Emergency department-based intervention with adolescent substance users: 12-month outcomes. *Drug Alcohol Dependence*, 79, 359-363.
- Taj, N., Devera-Sales, A., & Vinson, D. C. (1998). Screening for problem drinking: Does a single question work? *Journal of Family Practice*, 46(4), 328-335.
- Tenth Special Report to the U.S. Congress on Alcohol and Health. (2000). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism (NIAAA); NIH publication no. 00-1583.
- Terrell F., Zatzick D.F., Jurkovich G.J., Rivara F.P., Donovan D.M., Dunn C.W. et al. (2008). Nationwide survey of alcohol screening and brief intervention practices at US Level I trauma centers. (eng), *Journal of The American College of Surgeons*, 207 (5), 630-8.
- Timko, C., Moos, R. H., Finney, J. W., Moos, B. S., & Kaplowitz, M. S. (1999). Long-term treatment careers and outcomes of previously untreated alcoholics. *Journal of Studies on Alcohol*, 60, 437-447.
- Thomas B. (2004). Training Medical Providers to Conduct Alcohol Screening and Brief Interventions. *Substance Abuse*, 25 (1), 17-26.
- Tollison S.J., Lee C.M., Neighbors C., Neil T.A., Olson N.D., Larimer M.E. (2008). Questions and reflections: The use of motivational interviewing microskills in a peer-led brief alcohol intervention for college students. (eng), *Behavior Therapy*, 39 (2), 183-94.
- Toumbourou, J.W.; Stockwell, T.; Neighbors, C.; Marlatt, G.A.; Sturge, J.; Rehm, J. (2007). Interventions to reduce harm associated with adolescent substance use. *The Lancet (ScienceDirect)*, 69 (9570), 1391-1401.

- Turrisi, R., Larimer, M., Mallett, K., Kilmer, J., Ray, A., Mastroleo, N. et al. (2009), A Randomized Clinical Trial Evaluating a Combined Alcohol Intervention for High-Risk College Students. *Journal of Studies on Alcohol and Drugs*, 70 (4), 555-567.
- Unutzer, J., Katon, W., Callahan, C. M., et al. (2002). Collaborative care management of late-life depression in the primary care setting: a randomized controlled trial. *JAMA*, 2836-45.
- U.S. Preventive Services Task Force. *Screening for Depression*. (2002, May) Agency for Healthcare Research and Quality, Rockville, MD. <http://www.ahrq.gov/clinic/3rduspstf/depression/>
- U.S. Preventive Services Task Force (USPSTF). (2004). Screening and behavioral counseling interventions in primary care to reduce alcohol misuse: Recommendation statement. *Annals of Internal Medicine*, 140(7), 554-556.
- U.S. Preventive Services Task Force. (2009). Screening and treatment for major depressive disorder in children and adolescents: US preventative task force recommendation statement. *Pediatrics*, 123(4), 1223-1228
- [VanDeMark NR, Burrell NR, Lamendola WF, Hoich CA, Berg NP, Medina E.](#) (2010). An exploratory study of engagement in a technology-supported substance abuse intervention. [Subst Abuse Treat Prev Policy](#), Jun 8;5:10.
- Walker, EA, Torkelson N, Katon, WJ, & Koss, MP (1993). The prevalence rate of sexual trauma in a primary care clinic. *Journal of the American Board of Family Practice*, Sep-Oct; 6(5): 465-71
- Whiteside, U., Cronce, J. M., Pedersen, E. R., & Larimer, M. E. (2010). Brief motivational feedback for college students and adolescents: a harm reduction approach. *Journal of Clinical Psychology*, 66(2), 150–163.
- WHO ASSIST Working Group. (2002). The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): Development, reliability and feasibility. *Addiction*, 97, 1183–1194.
- Wilk, A.I., Jensen, N.M., and Havighurst, T.C. (1997). Meta-analysis of randomized control trails addressing brief interventions in heavy alcohol drinkers. *Journal of General Medicine*, 12 (5), 274-283.
- [Williams EC, Lapham G, Achtmeyer CE, Volpp B, Kivlahan DR, Bradley KA.](#) (2010). Use of an electronic clinical reminder for brief alcohol counseling for unhealthy alcohol use. [J Gen Intern Med](#). Jan; 25 Suppl 1:11-7.
- World Health Organization. (2008.) The effectiveness of a brief intervention for illicit drugs linked to the alcohol, smoking, and substance involvement screening test (ASSIST) in

primary health care settings: a technical report of phase III findings of the WHO ASSIST Randomized control trial. Retrieved June 16, 2010, from http://www.who.int/substance_abuse/activities/assist_technicalreport_phase3_final.pdf.

Zarkin, G.A., Bray, J.W., Davis, K.L., Babor, T.F., Higgins-Biddle, J.C. (2003). The costs of screening and brief intervention for risky alcohol use. (eng), *Journal of Studies on Alcohol*, 64 (6), 849-57.